arXiv:1006.0604v3 [math.DS] 2 Jun 2011

An example of unbounded chaos

Bau-Sen Du
Institute of Mathematics
Academia Sinica
Taipei 10617, Taiwan
dubs@math.sinica.edu.tw

Abstract

Let $\phi(x) = |1 - \frac{1}{x}|$ for all $x > 0$. Then we extend $\phi(x)$ in the usual way to become a continuous map from the compact topological (but not metric) space $[0, \infty]$ onto itself which also maps the set of irrational points in $(0, \infty)$ onto itself. In this note, we show that (1) on $[0, \infty]$, $\phi(x)$ is topologically mixing, has dense irrational periodic points, and has topological entropy $\log \lambda$, where λ is the unique positive zero of the polynomial $x^3 - 2x - 1$; (2) $\phi(x)$ has bounded uncountable *invariant* 2-scrambled sets of irrational points in $(0, 3)$; (3) for any countably infinite set X of points (rational or irrational) in $(0, \infty)$, there exists a dense unbounded uncountable *invariant* ∞-scrambled set Y of irrational transitive points in $(0, \infty)$ such that, for any $x \in X$ and any $y \in Y$, we have $\limsup_{n\to\infty} |\phi^n(x) - \phi^n(y)| = \infty$ and $\liminf_{n\to\infty} |\phi^n(x) - \phi^n(y)| = 0$. This demonstrates the true nature of chaos for $\phi(x)$ (see [2, 6, 7]).

Keywords: invariant scrambled sets, topological entropy, topological mixing, transitive points
AMS Subject Classification: 37D45, 37E05

1. Introduction

Let I denote an interval in the real line and let f be a continuous map from I into itself. It is well-known that if f has a periodic point of period not an integral power of 2 then there exist a positive number δ and an uncountable set S such that

$$\text{for any } x \neq y \text{ in } S,\ \limsup_{n\to\infty} |f^n(x) - f^n(y)| \geq \delta \ \text{ and } \ \liminf_{n\to\infty} |f^n(x) - f^n(y)| = 0. \qquad (*)$$

Such a set S is called a δ-scrambled set of f. When the inequality in (*) holds for all positive numbers δ (we follow the convention that $|\pm\infty \pm$ any real number$| = \infty$ and $\infty >$ any real number), we call such set S an ∞-scrambled set. Can this scrambled set S be chosen to be invariant under f? That is, can $f(S) \subset S$? The answer is no in general. This is because if c is a point in I such that $\liminf_{n\to\infty} |f^n(c) - f^n(f(c))| = 0$, then the ω-limit set $\omega(f, c)$ of

c must contain a fixed point of f, where the ω-limit set $\omega(f, c)$ of a point c is the set of all points x with the property that there is an increasing sequence $< n_i >$ of positive integers such that $\lim_{i \to \infty} f^{n_i}(c) = x$. This fact is useful in constructing examples of maps without *invariant* scrambled sets. For example, let g be the continuous map from $[0, 1]$ onto itself such that (i) $g(0) = 1$, $g(1) = 1/2$, and $g(1/2) = 0$; (ii) $g(1/6) = 1/3$ and $g(1/3) = 1/6$; and (iii) g is linear on each of the intervals $[0, 1/6]$, $[1/6, 1/3]$, $[1/3, 1/2]$ and $[1/2, 1]$. Then the point $x = 1/4$ is the unique fixed point of g and every point in $(1/6, 1/4) \cup (1/4, 1/3)$ is a period-2 point of g. Therefore, g has no *invariant* scrambled sets although g has the period-3 orbit $\{0, 1/2, 1\}$. On the other hand, if f is turbulent, i.e., if there exist two compact subintervals I_0 and I_1 of I with at most one point in common such that $f(I_0) \cap f(I_1) \supset I_0 \cup I_1$, then we can find such an *invariant* scrambled set [1].



Figure 1: The graph of the map $\phi(x) = |1 - \frac{1}{x}|, x > 0$.

In studying the periodicity and convergence of the second-order difference equation $x_{n+1} = |x_n - x_{n-1}|$, Sedaghat [3, 4] introduced the following interval map, see Figure 1,

$$\phi(x) = \left|1 - \frac{1}{x}\right| \text{ defined on } (0, \infty)$$

and showed, among other things, that $\phi(x)$ has uncountable scrambled sets. Following the work of Sedaghat [3, 4], we may ask the following two questions:

(1) Does $\phi(x)$ have bounded uncountable *invariant* scrambled sets ?

(2) Does $\phi(x)$ have unbounded uncountable *invariant* scrambled sets ?

In this note, we shall answer both questions afirmatively. We shall also show that for any countably infinite set X of points (rational or irrational) in $(0, \infty)$, there exists a dense unbounded uncountable invariant ∞-scrambled set Y of irrational transitive points (transitive points are points with dense orbits) in $(0, \infty)$ such that for any $x \in X$ and any $y \in Y$, we have $\limsup_{n \to \infty} |\phi^n(x) - \phi^n(y)| = \infty$ and $\liminf_{n \to \infty} |\phi^n(x) - \phi^n(y)| = 0$. For these purposes, we shall use symbolic dynamics.

2. Symbolic dynamics

Let $\Sigma_2 = \{\beta : \beta = \beta_0\beta_1 \cdots, \text{ where } \beta_i = 0 \,\text{or}\, 1\}$ be the metric space with metric d defined by

$$d(\beta_0\beta_1 \cdots, \gamma_0\gamma_1 \cdots) = \sum_{i=0}^{\infty} \frac{|\beta_i - \gamma_i|}{2^{i+1}}.$$

Let σ be the shift map defined by $\sigma(\beta_0\beta_1\beta_2 \cdots) = \beta_1\beta_2 \cdots$. Then σ is a continuous, two-to-one map from Σ_2 onto itself. In the sequel, for any finite sequences $\beta_0\beta_1\beta_2 \cdots \beta_k$, $k \geq 2$, of 0's and 1's, we also define $\sigma(\beta_0\beta_1\beta_2 \cdots \beta_k) = \beta_1\beta_2 \cdots \beta_k$.



Figure 2: Recursive procedures to obtain $I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i-1}\gamma_{n_i}}$'s.

Let $\phi(x) = |1 - 1/x|$ for all $x > 0$. We now compactify the space $[0, \infty)$ by including the symbol ∞ and define $\phi(0) = \infty$ and $\phi(\infty) = 1$. Then ϕ is a continuous map from the compact topological (but not metric) space $[0, \infty]$ onto itself. Let $I_0 = [0, 1]$ and $I_1 = [1, \infty]$. Then $\phi(I_0) = [0, \infty] = I_0 \cup I_1$ and $\phi(I_1) = [0, 1] = I_0$. Since $\phi(I_0) = I_0 \cup I_1$ and ϕ is strictly decreasing on I_0, we can split I_0 up into two compact subintervals I_{00} and I_{01} with one point in common such that $\phi(I_{0i}) = I_i = I_{\sigma(0i)}$, $i = 0, 1$. Since $\phi(I_1) = I_0$ which is disjoint from the interior of I_1, we cannot have two proper compact subintervals I_{10} and I_{11} of I_1 such that $\phi(I_{1i}) = I_i$, $i = 0, 1$. However, since $\phi(I_1) = I_0 = I_{00} \cup I_{01}$ and ϕ is strictly increasing on I_1, we can split I_1 up into two compact subintervals I_{100} and I_{101} with one point in common such that $\phi(I_{10i}) = I_{0i} = I_{\sigma(10i)}$, $i = 0, 1$. Therefore, although we *cannot* split I_1 up into two compact subintervals I_{10} and I_{11} with one point in common such that $\phi(I_{1i}) = I_i, i = 0, 1$, we can split I_1 up into two compact subintervals I_{100} and I_{101} (subscripts obtained by inserting the number 1 before each of 00 and 01) with one point in common such that $\phi(I_{10i}) = I_{0i}, i = 0, 1$. Similarly, we can continue with the above procedures (see Figure 2) as follows:

(1) since ϕ is strictly decreasing on $I_0 = [0,1]$ and $\phi(I_0) = [0,\infty] = I_{100} \cup I_{101} \cup I_{00} \cup I_{01}$ (in that order which is obtained from Steps 2 and 1 or, $(2n)^{th}$ and $(2n-1)^{st}$ steps in Figure 2 from right to left), we can split I_0 up, from left to right, into four compact subintervals with pairwise disjoint interiors, $I_{0100}, I_{0101}, I_{000}, I_{001}$ (obtained by inserting a 0 before the "previous" four subscripts) such that $I_{\gamma_0\gamma_1\cdots\gamma_{n_i}} \subset I_{\gamma_0\gamma_1\cdots\gamma_{n_i-1}}$ and $\phi(I_{\gamma_0\gamma_1\cdots\gamma_{n_i}}) = I_{\gamma_1\gamma_2\cdots\gamma_{n_i}}$ wherever defined, and the union of all these intervals is $[0,1]$,

(2) since ϕ is strictly increasing on $I_1 = [1,\infty]$ and $\phi(I_1) = [0,1] = I_{0100} \cup I_{0101} \cup I_{000} \cup I_{001}$ (in that order which is obtained from Procedure 1, i.e., Step 3 or $(2n+1)^{st}$ step in Figure 2 from left to right), we can split I_1 up, from left to right, into four compact subintervals with pairwise disjoint interiors, $I_{10100}, I_{10101}, I_{1000}, I_{1001}$ (obtained by inserting a 1 before the "previous" four subscripts) such that $I_{\gamma_0\gamma_1\cdots\gamma_{n_i}} \subset I_{\gamma_0\gamma_1\cdots\gamma_{n_i-1}}$ and $\phi(I_{\gamma_0\gamma_1\cdots\gamma_{n_i}}) = I_{\gamma_1\gamma_2\cdots\gamma_{n_i}}$ wherever defined, and the union of all these intervals is $[1,\infty]$,

(3) repeat Procedure 1 first and then Procedure 2 indefinitely.

Consequently, $I_0 = [0,1], I_1 = [1,\infty], I_{00} = [1/2,1], I_{01} = [0,1/2]$ and, for any finite sequence $\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i} = \gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i-2}00$ or $\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i-2}01$, $n_i \geq 2$, of 0's and 1's such that if $\gamma_j = 1$ for some $0 \leq j \leq n_i - 2$ then $\gamma_{j+1} = 0$, the compact interval $I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is defined so that $I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}} \subset I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i-1}}$ and $\phi(I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}}) = I_{\gamma_1\gamma_2\cdots\gamma_{n_i}}$. Note that, in Figure 2, if $I_{0\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is defined at $(2n-1)^{st}$ step, then $I_{10\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is defined at $(2n)^{th}$ step, and $I_{00\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is defined at $(2n+2)^{nd}$ step, while if $I_{1\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is defined at $(2n)^{th}$ step (and so $\gamma_1 = 0$), then $I_{01\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is defined at $(2n+1)^{st}$ step. Furthermore, at the $(2n-1)^{st}$ $((2n)^{th}$ respectively) step, the interval $[0,1]$ $([1,\infty]$ respectively) is split up into the union of 2^n compact subintervals with pairwise disjoint interiors and each of these 2^n intervals is split up at the next step into the union of two compact subintervals with disjoint interiors. Let $\Gamma = \{\gamma = \gamma_0\gamma_1\gamma_2\cdots \in \Sigma_2 : \text{if } \gamma_j = 1 \text{ then } \gamma_{j+1} = 0\}$. Then Γ is a subshift of finite type and, for any $\gamma = \gamma_0\gamma_1\gamma_2\cdots \neq \overline{100}$ in Γ ($\overline{100}$ stands for the sequence of repeating 100 in Γ and note that $I_{\overline{100}} = \{\infty\}$), the sequence $< I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}} >_{i\geq 1}$, wherever $I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is defined, is a nested sequence of compact intervals in $[0,\infty)$. Therefore, the set $I_\gamma = \cap_{i\geq 0} I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is either a nontrivial compact interval or consists of exactly one point and $\phi(I_\gamma) = I_{\sigma\gamma}$ for each $\gamma \in \Gamma$. However, since each rational point in $[0,\infty)$ is mapped to the point 0 after a finite number of iterations of ϕ [3], it is clear that each I_γ actually consists of exactly one point. *In the sequel*, we write this point as x_γ. So, $I_\gamma = \{x_\gamma\}$. Since $\phi(I_\gamma) = I_{\sigma\gamma}$ for each $\gamma \in \Gamma$, we obtain that if $\sigma^m(\gamma) = \gamma$ for some $\gamma \in \Gamma$, then $\phi^m(x_\gamma) = x_\gamma$. We note in passing that this correspondence $\gamma \to x_\gamma$ from Γ to $[0,\infty]$ is not one-to-one, but is *onto*. For each rational number $s > 0$, there are exactly two distinct $\gamma^{(1)}, \gamma^{(2)}$ in Γ such that $I_{\gamma^{(1)}} = I_{\gamma^{(2)}} = \{s\}$. For example, $I_{0\overline{010}} = \{1\} = I_{1\overline{010}}$. However, we don't need this information later on.

We remark that, since $\phi(x)$ is a very special map, we can actually compute these compact intervals $I_{\gamma_0\gamma_1\cdots\gamma_{n_i}}$ explicitly. They are related to Farey sequences in the way that, if $I_0 = [0,1] = [\frac{0}{1}, \frac{1}{1}], I_1 = [1,\infty] = [\frac{1}{1}, \frac{1}{0}]$, and if $a \geq 1, b \geq 0, c \geq 0, d \geq 1$ are integers such that $[\frac{b}{a}, \frac{d}{c}]$ (in lowest terms) is one of these compact intervals $I_{\gamma_0\gamma_1\cdots\gamma_{n_i}}$ in $[0,\infty]$ obtained at the n^{th} step, then $[\frac{b}{a}, \frac{b+d}{a+c}]$ and $[\frac{b+d}{a+c}, \frac{d}{c}]$ are two adjacent compact intervals in $[0,\infty]$ obtained at $(n+1)^{st}$ step (see Figure 2). However, we don't need this information to achieve our goal.

3. The extended map $\phi(x)$ is topologically mixing, has dense transitive irrational points and has dense irrational periodic points in $[0, \infty]$

Let V be a nonempty open interval in $(0, \infty)$, then there exist an irrational point w in V and an element $\gamma = \gamma_0\gamma_1\gamma_2\cdots$ in Γ such that $I_\gamma = \{w\}$. Since there exists a strictly increasing sequence $< n_i >$ of positive integers such that $V \supset \{w\} = I_\gamma = \cap_{i\geq 0} I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}}$, there is an integer $k \geq 5$ such that $w \in I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_k}} \subset V$. Consequently, $I_0 = \phi^{n_k+1}(I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_k}00}) \subset \phi^{n_k+1}(V)$ and so $[0, \infty] \supset \phi^{n_k+2}(V) \supset \phi(I_0) = [0, \infty]$. Thus, $\phi^{n_k+2}(V) = [0, \infty]$. This implies trivially that the extended map $\phi(x)$ is topologically mixing on $[0, \infty]$. Later in section 6, we shall see that $\phi(x)$ has dense transitive irrational points in $[0, \infty]$. Finally, since each rational point in $(0, \infty)$ is mapped into the period-3 orbit $\{1, 0, \infty\}$ under a finite number of iterations of ϕ and since $\{1\} = I_{\overline{001}} = I_{101\overline{001}}$, the unique point p in the set $I_{\overline{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_k}000}} \subset I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_k}000} \subset I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_k}} \subset V$ is an irrational periodic point of $\phi(x)$ in V. This shows that the irrational periodic points of $\phi(x)$ are dense in $(0, \infty)$.

4. The topological entropy of the extended map $\phi(x)$ on $[0, \infty]$

Since the extended map $\phi(x)$ is continuous on the compact topological space $[0, \infty]$, we can compute its topological entropy. We first define the continuous piecewise linear map $f : [0, 1] \to [0, 1]$ by putting $f(x) = 1 - 2x$ for $0 \leq x \leq 1/2$ and $f(x) = x - 1/2$ for $1/2 \leq x \leq 1$. Then $\{0, \frac{1}{2}, 1\}$ is a period-3 orbit of f and it is well-known that the topological entropy of f is $\log\lambda$, where λ is the unique positive zero of the polynomial $x^3 - 2x - 1$. In this section, we shall show that ϕ is topologically conjugate to the map f and so ϕ also has entropy $\log\lambda$. For this purpose, we define modified Farey sequences on $[0, \infty]$ inductively as follows: $F_0 = \{\frac{0}{1}, \frac{1}{0}\}$ and if $\frac{a}{b}$ and $\frac{c}{d}$ (both are in lowest terms) are two consecutive fractions of F_n, then $\frac{a}{b}, \frac{a+c}{b+d}, \frac{c}{d}$ are consecutive fractions of F_{n+1}. The first four modified Farey sequences are

$$F_0 = \{\frac{0}{1}, \frac{1}{0}\}; \quad F_1 = \{\frac{0}{1}, \frac{1}{1}, \frac{1}{0}\}; \quad F_2 = \{\frac{0}{1}, \frac{1}{2}, \frac{1}{1}, \frac{2}{1}, \frac{1}{0}\}; \quad F_3 = \{\frac{0}{1}, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \frac{1}{1}, \frac{3}{2}, \frac{2}{1}, \frac{3}{1}, \frac{1}{0}\}.$$

Furthermore, if we write, for each integer $n \geq 1$, $F_n = \{0/1 = a_{0,n}/b_{0,n},\ a_{1,n}/b_{1,n},\ a_{2,n}/b_{2,n}, \cdots, a_{2^{n-1},n}/b_{2^{n-1},n} = 1/1, \cdots, a_{2^n,n}/b_{2^n,n} = 1/0\}$, then it is easy to check that

(a) $a_{i,n}/b_{i,n} = b_{2^n-i,n}/a_{2^n-i,n}$ for all $0 \leq i \leq 2^{n-1}$;

(b) $a_{i,n}/b_{i,n} + a_{2^{n-1}-i,n}/b_{2^{n-1}-i,n} = 1$ for all $0 \leq i \leq 2^{n-1}$;

(c) $\phi(a_{2^n+i,n}/b_{2^n+i,n}) = a_{i,n}/b_{i,n}$ for all $0 \leq i \leq 2^{n-1}$ (by (a) and (b));

(d) $\phi(a_{i,n+1}/b_{i,n+1}) =$ (by (a)) $b_{2^n-i,n+1}/a_{2^n-i,n+1} - 1 = a_{2^n-i,n}/b_{2^n-i,n}$ for all $0 \leq i \leq 2^n$;

Now for each positive integer n, we define a continuous piecewise linear map $h_n(x)$ from $[0, \infty]$ into $[0, 1]$ by putting $h_n(a_{i,n}/b_{i,n}) = i/2^n$ for all $0 \leq i \leq 2^n - 1$, $h_n(\infty) = h_n(a_{2^n,n}/b_{2^n,n}) = (2^n - 1)/2^n$, and "connecting the dots". It is clear that the sequence

$< h_n(x) >$ converges uniformly on the non-compact metric space $[0, \infty)$ to a strictly increasing continuous map $h(x)$ with $\lim_{x\to\infty} h(x) = 1$. Let $h(\infty) = 1$. Then $h(x)$ is a homeomorphism from $[0, \infty]$ onto $[0, 1]$ and it follows from (c) and (d) above that $h \circ \phi = f \circ h$ on each F_n and so on $[0, \infty]$ by continuity, where f is defined above. Since ϕ is topologically conjugate to f and f has topological entropy $\log \lambda$, where λ is the unique positive zero of the polynomial $x^3 - 2x - 1$, so has ϕ. We Remark that the map h can also be obtained as the modified Minkowski map $h : [0, \infty] \to [0, 1]$ defined [5] by putting $h(\frac{0}{1}) = 0, h(\frac{1}{0}) = 1$, and $h(\frac{a+c}{b+d}) = \frac{1}{2}[h(\frac{a}{b}) + h(\frac{c}{d})]$ whenever $\frac{a}{b}$ and $\frac{c}{d}$ (both are in lowest terms) are two consecutive fractions in the modified Farey sequences and $h(\frac{a}{b})$ and $h(\frac{c}{d})$ are already defined, and extending h continuously to the whole interval $[0, \infty]$.

Now since I_γ consists of exactly one point for each $\gamma = \gamma_0\gamma_1\gamma_2 \cdots \in \Gamma$ and $\phi(I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}}) = I_{\sigma(\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i})}$, wherever $I_{\gamma_0\gamma_1\gamma_2\cdots\gamma_{n_i}}$ is defined, it is easy to see that if γ and $\gamma(m)$, $m \ge 1$, are points in Γ such that $\lim_{m\to\infty} \gamma(m) = \gamma$, then $\lim_{m\to\infty} x_{\gamma(m)} = x_\gamma$. It is also easy to see that $I_{\bar{0}} = \{(\sqrt{5}-1)/2\}$ where $z = (\sqrt{5}-1)/2$ is the unique fixed point of $\phi(x)$, $I_{1\bar{0}} = \{(3+\sqrt{5})/2\}$, $I_{\overline{010}} = \{0\}$, $I_{\overline{001}} = \{1\}$, and $I_{\overline{100}} = \{\infty\}$. This fact will be needed below.

5. $\phi(x)$ has bounded uncountable invariant 2-scrambled sets in $[0, 3]$

The existence of bounded uncountable *invariant* δ-scrambled sets for some $\delta > 0$ follow easily from Theorem 3 of [1]. Here we use symbolic dynamics to give a different proof. For each $\beta = \beta_0\beta_1\beta_2 \cdots \in \Sigma_2$, let $\mu_\beta = (\mu_\beta)_0(\mu_\beta)_1(\mu_\beta)_2 \cdots$ be a point in Γ defined by

$$\mu_\beta = 0^{5!}\, A_\beta(5!)\, A_\beta(6!)\, A_\beta(7!) \,\cdots ,$$

where $0^2 = 00, 0^3 = 000$, etc., and for each $k \ge 5$, $A_\beta(k!) = (\mu_\beta)_{k!}(\mu_\beta)_{k!+1}(\mu_\beta)_{k!+2} \cdots (\mu_\beta)_{(k+1)!-1}$ is the concatenation of the following k strings of 0's and 1's, each of length $k!$,

$$010^{k!-2},\ 0\beta_0 0^{k!-2},\ 0\beta_1 0^{k!-2},\ 0\beta_2 0^{k!-2},\ \cdots ,\ 0\beta_{k-2}0^{k!-2}.$$

Let $W = \{\phi^n(x_{\mu_\beta}) : \beta \in \Sigma_2,\ n \ge 0\}$. Then it is clear that $W\,(\subset I_{000} \cup I_{001} \cup I_{0100} \cup I_{1000} = [0, 1/3] \cup [1/2, 1] \cup [2, 3]\,)$ is a bounded uncountable invariant subset of $[0, 3]$. If $\eta = \eta_0\eta_1\eta_2 \cdots$ and $\xi = \xi_0\xi_1\xi_2 \cdots$ are any two points (need not be distinct) in Σ_2 and if $i \ge 1$ and $m \ge 0$ are fixed integers, then, for each $k > i + m + 5$, we have

$$\sigma^{k!+1}(\mu_\eta) = 10^{k!-2} \cdots \quad \text{and} \quad \sigma^{k!+2}(\mu_\eta) = 0^{k!-2} \cdots ,$$

$$\sigma^{k!+1}(\sigma^i(\mu_\xi)) = 0^{k!-1-i} \cdots \quad \text{and} \quad \sigma^{k!+2}(\sigma^i(\mu_\xi)) = 0^{k!-2-i} \cdots ,$$

and

$$\sigma^{(m+2)k!+1}(\mu_\eta) = \eta_m 0^{k!-2} \cdots \quad \text{and} \quad \sigma^{(m+2)k!+1}(\mu_\xi) = \xi_m 0^{k!-2} \cdots .$$

As k tends to ∞, the sequence $< 10^{k!-2} \cdots >$ tends to the point $1\bar{0}$ and the sequence $< 0^{k!-2-i} \cdots >$ tends to the point $\bar{0}$. Since $\lim_{n\to\infty} I_{0^{k!-2-i}\cdots} = I_{\bar{0}} = \{(\sqrt{5} - 1)/2\} = \{x_{\bar{0}}\} = \{$the unique fixed point of $\phi\}$ and $\lim_{n\to\infty} I_{10^{k!-2}\cdots} = I_{1\bar{0}} = \{(3 + \sqrt{5})/2\} = \{x_{1\bar{0}}\} = \{$the unique inverse image in $(1, \infty)$ of the unique fixed point of $\phi\}$, we obtain

$$\limsup_{n\to\infty} |\phi^n(x_{\mu_\eta}) - \phi^n(\phi^i(x_{\mu_\xi}))| \ge |x_{1\bar{0}} - x_{\bar{0}}| = 2 \quad \text{and} \quad \liminf_{n\to\infty} |\phi^n(x_{\mu_\eta}) - \phi^n(\phi^i(x_{\mu_\xi}))| = 0.$$

Similarly, if $\eta_m \neq \xi_m$, then we have

$$\limsup_{n\to\infty} |\phi^n(x_{\mu_\eta}) - \phi^n(x_{\mu_\xi})| \geq 2 \quad \text{and} \quad \liminf_{n\to\infty} |\phi^n(x_{\mu_\eta}) - \phi^n(x_{\mu_\xi})| = 0.$$

Therefore, we have proved the following result:

Theorem 1. *Let* $\phi(x) = |1 - 1/x|$ *for all* $x > 0$. *Then* W ($\subset [0, 3]$) *is a bounded uncountable invariant 2-scrambled set for* $\phi(x)$.

6. $\phi(x)$ has dense unbounded uncountable invariant ∞-scrambled sets

Let $k \geq 5$ be a fixed integer. We call any finite sequence $\beta_0\beta_1\beta_2 \cdots \beta_k$ of 0's and 1's admissible if $\beta_i = 1$ and $i < k$ then $\beta_{i+1} = 0$. There are countably infinitely many such admissible finite sequences of 0's and 1's. Let $B_1, B_2, \cdots, B_n, \cdots$ be an enumeration of these admissible finite sequences with $B_n = \beta_{n,0}\beta_{n,1}\beta_{n,2}\cdots\beta_{n,k_n-1}$, $n \geq 1$. Let $m_1, m_2, \cdots$ be a strictly increasing sequence of positive integers such that $(m_i)! > (m_{i-1})! + k_{i-1} + 1$ for all $i \geq 2$. Let $\alpha = \alpha_0\alpha_1\alpha_2\cdots$ be a point in Σ_2 defined by putting $\alpha_{(m_i)!}\alpha_{(m_i)!+1}\alpha_{(m_i)!+2}\cdots\alpha_{(m_i)!+k_i-1} = B_i$ for all $i \geq 1$ and $\alpha_\ell = 0$ elsewhere. Then it is clear that α is a *totally transitive point* of σ in $\Gamma = \{\gamma = \gamma_0\gamma_1\gamma_2\cdots \in \Sigma_2 : \text{if } \gamma_i = 1 \text{ then } \gamma_{i+1} = 0\}$, i.e., for each integer $n \geq 1$, the orbit of the point α with respect to the map σ^n is dense in Γ.

Let $\alpha = \alpha_0\alpha_1\alpha_2\cdots$ be a *totally transitive point* of σ in Γ (and so if $I_\alpha = \{x_\alpha\}$ then x_α is a *totally transitive point* of ϕ in $(0, \infty)$). Let $X = \{x(1), x(2), x(3), \cdots\}$ be any countably infinite subset of *irrational* points in $(0, \infty)$. For each integer $m \geq 1$, we can write $\{x(m)\} = I_{\gamma^{(m)}}$ for an unique $\gamma^{(m)} = \gamma_0^{(m)}\gamma_1^{(m)}\gamma_2^{(m)}\cdots$ in Γ. For any integers $m \geq 1$ and $5 \leq i < j$ such that $j - i + 1$ is a multiple of 3, let

$$C(x(m),\, i : j) = \gamma_i^{(m)}\gamma_{i+1}^{(m)}\cdots\gamma_{j-1}^{(m)}\, 0 \quad \text{and}$$

$$C^*(x(m),\, i : j) = \xi_i^{(m)}\xi_{i+1}^{(m)}\xi_{i+2}^{(m)}\cdots\xi_j^{(m)} = \begin{cases} 0\,100\,100\,100\,\cdots 100\,10, & \text{if } \gamma_i^{(m)} = 1, \\ 100\,100\,100\,\cdots 100, & \text{otherwise.} \end{cases}$$

Note that, in $C(x(m),\, i : j)$ and in $C^*(x(m),\, i : j)$, the last element is 0 so when we concatenate C's with C's or with C^*'s, we won't have the finite string $\cdots 11 \cdots$. For simplicity, for $b = 0$ or 1, we write $(b00)^1 = b00$, $(b00)^2 = b00\,b00$, and so on.

For any $\beta = \beta_0\beta_1\beta_2\cdots$ in Σ_2, define a new point $\tau_\beta = \tau_\beta(\alpha, X)$ in Γ as follows and let

$$Y = \{\phi^n(x_{\tau_\beta}) : \beta \in \Sigma_2,\, n \geq 0\},$$

where $\tau_\beta = (\tau_\beta)_0(\tau_\beta)_1(\tau_\beta)_2\cdots = \alpha_0\alpha_1\alpha_2\cdots\alpha_{5!-2}0\ A_\beta(5!)\ A_\beta(6!)\ A_\beta(7!)\ \cdots$, and, for each $k \geq 5$, $A_\beta(k!) = (\tau_\beta)_{k!}(\tau_\beta)_{k!+1}(\tau_\beta)_{k!+2}\cdots(\tau_\beta)_{(k+1)!-1}$ is the concatenation of the following k strings of 0's and 1's, each of length $k!$,

$\alpha_0\alpha_1\alpha_2\cdots\alpha_{k!-1}$

$0^{k!/4}(100)^{k!/12}(001)^{k!/12}(010)^{k!/12}$

$$(\beta_0 00)^{(k-1)!/3}(\beta_1 00)^{(k-1)!/3}\cdots(\beta_{k-1}00)^{(k-1)!/3}$$

$$B(x(1),4k!) \qquad B(x(2),5k!) \qquad B(x(3),6k!) \qquad \cdots \qquad B(x(k-3),k\cdot k!),$$

where, for $1 \le i \le k-3$, $B(x(i),(3+i)k!)$ is the concatenation of the following $2k$ strings of 0's and 1's, each of length $\frac{1}{2}(k-1)!$,

$$C(x(i),(3+i)k! : (3+i)k! + [\tfrac{1}{2}(k-1)!-1])$$

$$\cdots$$

$$C(x(i),(3+i)k! + (j-1)[\tfrac{1}{2}(k-1)!-1] : (3+i)k! + j[\tfrac{1}{2}(k-1)!-1])$$

$$\cdots$$

$$C(x(i),(3+i)k! + (k-1)[\tfrac{1}{2}(k-1)!-1] : (3+i)k! + k[\tfrac{1}{2}(k-1)!-1])$$

$$C^*(x(i),(3+i)k! + \tfrac{1}{2}k! : (3+i)k! + \tfrac{1}{2}k! + [\tfrac{1}{2}(k-1)!-1)])$$

$$\cdots$$

$$C^*(x(i),(3+i)k! + \tfrac{1}{2}k! + (j-1)[\tfrac{1}{2}(k-1)!-1] : (3+i)k! + \tfrac{1}{2}k! + j[\tfrac{1}{2}(k-1)!-1])$$

$$\cdots$$

$$C^*(x(i),(3+i)k! + \tfrac{1}{2}k! + (k-1)[\tfrac{1}{2}(k-1)!-1] : (3+i)k! + \tfrac{1}{2}k! + k[\tfrac{1}{2}(k-1)!-1]).$$

In the following, we shall use the fact that $I_{\overline{100}} = \{\infty\}$, $I_{\overline{001}} = \{1\}$, $I_{\overline{010}} = \{0\}$, and if γ and $\gamma^{(n)}$, $n \ge 1$, are points in Γ such that $\lim_{n\to\infty} \gamma^{(n)} = \gamma$ then $\lim_{n\to\infty} \phi(x_{\gamma^{(n)}}) = x_\gamma$, where $I_{\gamma^{(n)}} = \{x_{\gamma^{(n)}}\}$ and $I_\gamma = \{x_\gamma\}$. Now in the expansion of τ_β, $\beta \in \Sigma_2$,

(1) there are infinitely many strings $\alpha_0\alpha_1\alpha_2\cdots\alpha_{k!-1}$, $k \ge m$, which imply, for each $i \ge 1$, the denseness of the orbit (under ϕ^i) of the point x_{τ_β} in $(0,\infty)$,

(2) there are infinitely many strings $(\beta_0 00)^{(k-1)!/3}(\beta_1 00)^{(k-1)!/3}\cdots(\beta_{k-1}00)^{(k-1)!/3}$, $k \ge 5$ which imply that

$$\limsup_{n\to\infty} |\phi^n(x_{\tau_\beta}) - \phi^n(x_{\tau_\eta})| = \infty \text{ for } \beta \ne \eta \text{ in } \Sigma_2,$$

(3) for any $i \ge 1$, there are infinitely many strings $0^{k!/4}(100)^{k!/12}(001)^{k!/12}(010)^{k!/12}$, $k \ge i+5$, containing

$$0^i(100)^{k!/12}(001)^{k!/12}(010)^{k!/12}$$

which imply that, for any integer $i \ge 1$,

$$\limsup_{n\to\infty} |\phi^n(x_{\tau_\beta}) - \phi^n(\phi^i(x_{\tau_\eta}))| = \infty \text{ for any } \beta \text{ and } \eta \text{ in } \Sigma_2,$$

and since this string also contains long strings of 0's, i.e., $0^{k!/4}$, $k \ge 5$, we obtain

$$\liminf_{n\to\infty} |\phi^n(x_{\tau_\beta}) - \phi^n(\phi^i(x_{\tau_\eta}))| = 0 \;\text{ for any }\; \beta, \eta \;\text{ in }\; \Sigma_2 \;\text{ and }\; i \ge 0,$$

(4) for any positive integers i and j, there are infinitely many strings

$$C(x(i), (3+i)k! + (j-1)[\frac{1}{2}(k-1)! - 1] : (3+i)k! + j[\frac{1}{2}(k-1)! - 1]),\; k > j,$$

which imply that

$$\liminf_{n\to\infty} |\phi^n(x(i)) - \phi^n(\phi^{j-1}(x_{\tau_\beta}))| = 0 \;\text{ for all }\; \beta \;\text{ in }\; \Sigma_2 \;\text{ and }\; j \ge 1,$$

(5) for any positive integers i and j, there are infinitely many strings

$$C^*(x(i), (3+i)k! + \frac{1}{2}k! + (j-1)[\frac{1}{2}(k-1)! - 1] : (3+i)k! + \frac{1}{2}k! + j[\frac{1}{2}(k-1)! - 1]),\; k > j,$$

which, since $x_{0\gamma_1\gamma_2\cdots} \le 1 \le x_{1\gamma_1\gamma_2\cdots}$ and $\lim_{k\to\infty} I_{(100)^k} = \{x_{\overline{100}} = \infty\}$, imply that

$$\limsup_{n\to\infty} |\phi^n(x(i)) - \phi^n(\phi^{j-1}(x_{\tau_\eta}))| = \infty \;\text{ for all }\; \beta \;\text{ in }\; \Sigma_2 \;\text{ and }\; j \ge 1.$$

Now, if r is a rational number in $[0, \infty)$, then, by [3], for some integer $m \ge 0$, $\phi^m(r) = 0$ which is a period 3 point of ϕ. For each β in Σ_2, the iterates of the point x_{τ_β} approach the fixed point $z = (\sqrt{5}-1)/2$ infinitely often and stay close to it for a while each time. This implies that $\limsup_{n\to\infty} |\phi^n(r) - \phi^n(x_{\tau_\beta})| = \infty$. Furthermore, since, for all $k > 12$,

$$\sigma^{3(k!/3+k!/12)}(\tau_\beta) = \sigma^{k!+k!/4}(\tau_\beta) = (100)^{k!/12}(001)^{k!/12}(010)^{k!/12}\cdots,$$

$$\sigma^{3(k!/3+k!/12+k!/36)}(\tau_\beta) = \sigma^{k!+k!/4+k!/12}(\tau_\beta) = (001)^{k!/12}(010)^{k!/12}\cdots,$$

and

$$\sigma^{3(k!/3+k!/12+k!/36+k!/36)}(\tau_\beta) = \sigma^{k!+k!/4+k!/12+k!/12}(\tau_\beta) = (010)^{k!/12}\cdots,$$

the sequence $< \sigma^{3n}(\tau_\beta) >$ approaches each of the three points $\overline{100}$, $\overline{001}$ and $\overline{010}$ infinitely often and so the sequence $< \phi^{3n}(x_{\tau_\beta}) >$ approaches each of the three points $\infty, 1, 0$ infinitely often. Since, for some $m \ge 1$, $\phi^m(r) = 0$ and 0 is a period-3 point of $\phi(x)$, this implies that $\liminf_{n\to\infty} |\phi^n(r) - \phi^n(x_{\tau_\beta})| = 0$ (if $f^{3n}(r) = \infty$ and $\phi^{3n}(x_{\tau_\beta}) \approx \infty$ then $\phi^{3n+1}(r) = 1$ and $\phi^{3n+1}(x_{\tau_\beta}) \approx 1$). Therefore, we have shown the following result:

Theorem 2. *Let $\phi(x) = |1 - 1/x|$ for all $x > 0$. Then for any given countably infinite subset X of points (rational or irrational) in $(0, \infty)$, there exists a dense unbounded uncountable invariant ∞-scrambled set Y of totally transitive irrational points in $(0, \infty)$ such that, for any x in X and any y in Y, we have*

$$\limsup_{n\to\infty} |\phi^n(x) - \phi^n(y)| = \infty \quad \textit{and} \quad \liminf_{n\to\infty} |\phi^n(x) - \phi^n(y)| = 0.$$

In the above result, we consider $\phi(x)$ as a continuous map from the compact topological (but not metric) space $[0, \infty]$ onto itself under the convention that $|\frac{1}{0}| = \infty$, $\frac{1}{\pm\infty} = 0$, and $| \pm \infty \pm$ any real number$| = \infty$. It is easy to see that the dynamics of $\phi(x)$ on the rational points in $[0, \infty)$ are not interesting [3, 4] because they all go to the period-3 orbit $\{1, 0, \infty\}$ after a finite number of iterations. All interesting dynamics occur in the invariant set $\mathbb{R}_+ \setminus \mathbb{Q}_+$ of irrational points of $(0, \infty)$. Consequently, we can consider the dynamics of $\phi(x)$ on the set $\mathbb{R}_+ \setminus \mathbb{Q}_+$ which is a metric space under the usual distance metric. Within this context, Theorem 2 has a very important consequence: given any *irrational* point x in $\mathbb{R}_+ \setminus \mathbb{Q}_+$, then at just about everywhere in $\mathbb{R}_+ \setminus \mathbb{Q}_+$, whether it is close to x or far away from it we can always find an *irrational* point y (in the dense set Y) whose iterates satisfy $\limsup_{n\to\infty} |\phi^n(x) - \phi^n(y)| = \infty$ and $\liminf_{n\to\infty} |\phi^n(x) - \phi^n(y)| = 0$. This demonstrates the true nature of chaos [2, 6, 7], i.e., not only nearby points will separate (sensitivity) and converge infinitely often but even *far apart* points will also converge and separate infinitely often.

Acknowledgement

This work was partially supported by the National Science Council NSC 97-2119-M-001-004.

References

[1] B.-S. Du, *On the invariance of Li-Yorke chaos of interval maps,* J. Diff. Equ. Appl. 11 (2005), pp. 823-828.

[2] B.-S. Du, *On the nature of chaos,* arXiv:math/0602585 (2006).

[3] H. Sedaghat, *Periodicity and convergence for* $x_{n+1} = |x_n - x_{n-1}|$, J. Math. Anal. Appl. 291 (2004), pp. 31-39.

[4] H. Sedaghat, *The Li-Yorke theorem and infinite discontinuities,* J. Math. Anal. Appl. 296 (2004), pp. 538-540.

[5] R. Girgensohn, *Constructing singular functions via Farey fractions,* J. Math. Anal. Appl. 203 (1996), pp. 127-141.

[6] A. Vieru, *General definitions of chaos for continuous and discrete-time processes,* arXiv:0802.0677v3 (2008).

[7] A. Vieru, *About stable periodic helixes, L-iteration and chaos generated by unbounded functions,* arXiv:0802.1401v2 (2008).